MULTIMEDIA UNIVERSITY

STUDENT IDENTIFICATION NO

MULTIMEDIA UNIVERSITY

FINAL EXAMINATION

TRIMESTER 2, 2018/2019 SESSION

BMS2024 - ADVANCED MANAGERIAL STATISTICS

(All Sections / Groups)

04 MARCH 2019
9am – 11am
(2 Hours)

INSTRUCTIONS TO STUDENTS

1. This question paper consists of 14 pages excluding the cover page.
2. This question paper consists of **FOUR** structured questions. Attempt **ALL** questions.
3. Students are allowed to use non-programmable scientific calculators with no restrictions.
4. Statistical tables are attached at the end of the question paper.
5. Please use **pen** to write the answers.
6. Please write all your answers in the answer booklet provided.

QUESTION 1 [25 MARKS]

a) The University had received complaints from the students and faculty about lateness, the registrar at a large university is ready to undertake a study to determine whether the scheduled break between classes should be changed. Until now the registrar has believed that there should be 20 minutes gap between scheduled classes. State the null hypothesis, H_0, and the alternative hypothesis, H_1. [4 marks]

b) Bottles of water have a label stating that the volume is 12 oz. A consumer group suspects the bottles are under-filled and plans to conduct a test. Define type I and II error in this case?

[4 marks]

c) Trace metals in drinking water affect the flavor and an unusually high concentration can pose a health hazard. Ten pairs of data were taken measuring zinc concentration in bottom water and surface water. The data given in the table below. Does the data suggest that the true average concentration in the bottom water exceeds that of surface water? Perform the hypothesis testing and interpret your answer.

Let A be the Zinc Concentration in bottom water and;
Let B be the Zinc Concentration in surface water.

Location	1	2	3	4	5	6	7	8	9	10
A	0.430	0.266	0.567	0.531	0.707	0.716	0.651	0.589	0.469	0.723
B	0.415	0.238	0.390	0.410	0.605	0.609	0.632	0.523	0.411	0.612

[17 marks]

[Total : 25 Marks]

Continued...

QUESTION 2 [25 MARKS]

a) Explain the differences between parametric and nonparametric tests. Discuss in terms of their assumptions and the circumstances under which each should be applied and provide one example of such test.

[5 marks]

b) Does physical exercise alleviate depression? A random number of depressed people were selected to answer the survey. They were then allocated in three groups: no exercise, 20 minutes of jogging per day, or 60 minutes of jogging per day. At the end of the month, each participant was asked to rate how depressed he or she now feel, on a likert scale that runs from 1 ("totally miserable") through to 100 (ecstatically happy"). Is there evidence of significance difference in the median result between those participants? Be sure to show all steps and workings.

[20 marks]

No exercise	Jogging for 20 minutes	Jogging for 60 minutes
23	22	59
26	27	66
51	39	38
49	29	49
58	46	56
37	48	60
29	49	56
44	65	62

[Total : 25 Marks]

Continued...

QUESTION 3 [25 Marks]

The US Census Bureau collects statistics comparing the various 50 states. Some of the data collected has been single out the first three variables; infant mortality rate per 1,000 live births), white (i.e. the percentage of the population that is white), and the violent crime rate (per 100,000 people). The data can be analysed in which the poverty rate (% of population below the poverty level) can be estimated as a multiple linear combination of the infant mortality rate, the percentage of the population that is white and the violent crime rate.

The summary output of the analysis is shown below:

ANOVA

	df	*SS*	*MS*	*F*	*Significance F*
Regression	3	142.5503596	47.51678652	7.786905323	$2.6621*10^{-4}$
Residual	46	280.6984404	6.102140009		
Total	49	423.2488			

	Coefficients	*Std Error*	*t Stat*	*P-value*
Intercept	0.43712519	3.98753369	0.109622953	
Infant Mort	1.279369653	0.300672909	4.255021369	0.000101628
White	0.036326923	0.033602532	1.08107696	0.285298153
Crime	0.001421499	0.002242102	0.634002914	0.529219218

a) State the multiple linear regression equation for the above data. [4 marks]

b) Interpret the slope coefficient for the infant mortality and crime rate relating to the poverty rate. [4 marks]

c) How strong is the model? Compute and interpret the value. [4 marks]

d) At the 5% level of significance, use the p-value approach to test the overall validity of the model. [4 marks]

e) At the 1% level of significance, use the p-value approach to test if each independent variable is significantly related towards poverty rate. [6 marks]

f) Predict what would be the poverty rate in a state with 5,600 live birth, 80% of white people and 250,000 crimes. [3 marks]

[Total : 25 Marks]

Continued...

QUESTION 4 [25 Marks]

Suppose the Company ABC wants to examine the safety of compact cars, midsize cars, and full-size cars. It collects a sample of three for each of the treatments (cars types). Using the hypothetical data provided below, test whether the mean pressure applied to the driver's head during a crash test is equal for each types of car. Use $\alpha = 5\%$. Assume that the dataset is normally distributed.

Summary Output

Groups	Count	Sum	Mean	Variance
Compact Cars	3	2000	666.6667	972.3333
Midsize Cars	3	1421	473.6667	2417.333
Full-size Cars	3	1342	447.3333	1737.333.

ANOVA

Source of Variation	SS	df	MS	F
Among Groups	86049.55556	2	**43024.78**	**25.17541**
Within Groups	10254	6	1709	
Total	96303.55556	8		

a) Show the working to get MSA = 43024.78 and F=25.17541. [5 marks]

b) At the 5% level of significance, is there evidence of a difference in the mean pressure applied to the driver's head during a crash test for each types of car? Conduct an appropriate statistical procedure. [8 marks]

c) Conduct the Tukey-Kramer post-hoc test to examine which types of car differ in mean pressure applied to the driver's head during a crash test. Use 10 % significance level. [12 marks]

[Total : 25 Marks]

End of Page

STATISTICAL FORMULAE

A. DESCRIPTIVE STATISTICS

Sample Mean = $\overline{X} = \dfrac{\sum_{i=1}^{n} X_i}{n}$ Sample Standard Deviation = s = $\sqrt{\dfrac{\sum_{i=1}^{n} X_i^2}{n-1} - \dfrac{\left(\sum_{i=1}^{n} X_i\right)^2}{n(n-1)}}$

where n = *number of observations*
X_i = *the* i^{th} *observation of the data*

B. HYPOTHESIS TESTING

Types of Error
Type I Error = α= P(Rejecting H_0 | H_0 is true) where, Confidence Interval = 1 - α
Type II Error = β= P(Not Rejecting H_0 | H_0 is false)

One Sample Mean Test	
σ Known	**σ Unknown**
$z = \dfrac{\overline{x} - \mu}{\sigma/\sqrt{n}}$	$t = \dfrac{\overline{x} - \mu}{s/\sqrt{n}}$

Two Sample Mean Test	
Comparing Means for Two Independent Populations	
[Standard Deviation (σ) Known] $z = \dfrac{(\overline{x_1} - \overline{x_2}) - (\mu_1 - \mu_2)}{\sqrt{\sigma_1^2/n_1 + \sigma_2^2/n_2}}$	**[Standard Deviation (σ) Not Known]** $t = \dfrac{(\overline{x_1} - \overline{x_2}) - (\mu_1 - \mu_2)}{\sqrt{S_p^2\left(1/n_1 + 1/n_2\right)}}$ where $s_p^2 = \dfrac{(n_1-1)S_1^2 + (n_2-1)S_2^2}{(n_1+n_2-2)}$

Two Sample Mean Test
Comparing Means for Two Paired Populations
$t = \dfrac{(\overline{D} - \mu_D)}{S_D/\sqrt{n}}$ where $\overline{D} = \dfrac{\sum_{i=1}^{n} D_i}{n}$ and $S_D = \sqrt{\dfrac{\sum_{i=1}^{n} D_i^2}{n-1} - \dfrac{\left(\sum_{i=1}^{n} D_i\right)^2}{n(n-1)}}$

Non-Parametric Analysis

Wilcoxon Rank Sum Test	Wilcoxon Signed Rank Sum Test
$Z = \frac{(T_1 - \mu_{T_1})}{\sigma_{T_1}}$ where	$Z = \frac{(T_+ - \mu_{T_+})}{\sigma_{T_+}}$ where
$\mu_{T1} = \frac{n_1(n+1)}{2}$ and	$\mu_{T+} = \frac{n(n+1)}{4}$ and
$\sigma_{T_1} = \sqrt{\frac{n_1 n_2 (n+1)}{12}}$ where $n = n_1 + n_2$	$\sigma_{T_+} = \sqrt{\frac{n(n+1)(2n+1)}{24}}$

Kruskal-Wallis Rank Test

$H = \left[\frac{12}{n(n+1)} \sum_{j=1}^{c} \frac{T_j^2}{n_j} \right] - 3(n+1)$ where the critical value is χ^2 with $df = c - 1$

Check ranking sum: $\sum T_j = n(n+1)/2$

Chi-Square Test

$$\chi^2 = \sum^{n} \frac{(O-E)^2}{E}$$

where O = Frequency of Observed Values and E = Frequency of Expected Values

with df = c – 1 where c = number of categories
or
with df = (r – 1)(c – 1) where r = number of rows and c = number of columns

C. ANALYSIS OF VARIANCE (ANOVA)

One-Way ANOVA

Source	Degrees of Freedom	Sum of Squares	Mean Squares	F-statistic
Among Groups	*c - 1*	SSA	MSA = SSA/*c-1*	MSA/MSW
Within Groups	*n - c*	SSW	MSW = SSW/*n-c*	
Total	*n - 1*	SST		

$$SST = \sum_{j=1}^{c}\sum_{i=1}^{n_j}\left(X_{ij} - \overline{\overline{X}}\right)^2 \text{ or alternative formula: } SST = \left(\sum_{j=1}^{c}\sum_{i=1}^{n_i} X_{ij}^2\right) - \frac{\left(\sum_{j=1}^{c}\sum_{i=1}^{n_i} X_{ij}\right)^2}{n}$$

$$SSA = \sum_{j=1}^{c} n_j\left(\overline{X}_j - \overline{\overline{X}}\right)^2 \text{ and } SSW = SST - SSA$$

where n = number of observations, c = number of groups and $\overline{\overline{X}}$ *= overall mean*

Tukey-Kramer Procedure

$$\text{Critical Range} = Q_U\sqrt{\frac{MSW}{2}\left[\frac{1}{n_i} + \frac{1}{n_j}\right]}$$

where Q_u *= the upper tail critical value from a Studentized Range Distribution having (c) degrees of freedom in the numerator and (n – c) degrees of freedom in the denominator at a given level of significance*

Two-Way ANOVA

Source	Degrees of Freedom	Sum of Squares	Mean Squares	F-statistic
A	$r-1$	SSA	MSA = SSA/$(r-1)$	MSA / MSE
B	$c-1$	SSB	MSB = SSB/$(c-1)$	MSB / MSE
AB	$(r-1)(c-1)$	SSAB	MSAB = SSAB/$(r-1)(c-1)$	MSAB / MSE
Error	$rc(n-1)$	SSE	MSE = SSE/$rc(n'-1)$	
Total	$n-1$	SST		

where,

Factor A levels are represented by the rows and Factor B levels are represented by the columns and
n = number of observations
c = number of columns
r = number of rows
n' = number of replicates

$$SST = \sum_{i=1}^{r}\sum_{j=1}^{c}\sum_{k=1}^{n'}\left(X_{ijk} - \overline{\overline{X}}\right)^2 \qquad SSA = cn'\sum_{i=1}^{r}\left(\overline{X}_i - \overline{\overline{X}}\right)^2$$

$$SSB = rn'\sum_{j=1}^{c}\left(\overline{X}j - \overline{\overline{X}}\right)^2 \quad \text{where } \overline{\overline{X}} = \textit{overall mean}$$

$$SSE = (n'-1)\left[S_1^2 + S_2^2 + \ldots\ldots + S_k^2\right] \quad \text{where } S_i^2 = \text{variance of each block}$$

D. REGRESSION ANALYSIS

Multiple Linear Regression

Population Model: $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \ldots\ldots + \beta_k X_k + \varepsilon$

Sample Model: $\hat{y} = b_0 + b_1 x_1 + b_2 x_2 + \ldots\ldots + b_k x_k + e$

Adjusted R-Square $= 1 - \left[\dfrac{(1-R^2)(n-1)}{(n-p-1)}\right]$ where p = number of independent/predictor variables

ANOVA Table for Regression

Source	Degrees of Freedom	Sum of Squares	Mean Squares
Regression	p	SSR	MSR = SSR/p
Error/Residual	$n-p-1$	SSE	MSE = SSE/$(n-p-1)$
Total	$n-1$	SST	

Test Statistic for Significance of the Overall Regression Model

F = MSR/MSE

Test Statistic for Significance of Each Predictor Variable

$t_i = \dfrac{b_i}{S_{b_i}}$ and the critical value $= \pm t_{\alpha/2,(n-p-1)}$

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

The function tabulated is $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-\frac{1}{2}t^2}\, dt$. $\Phi(x)$ is the probability that a random variable, normally distributed with zero mean and unit variance, will be less than or equal to x. When $x < 0$ use $\Phi(x) = 1 - \Phi(-x)$, as the normal distribution with zero mean and unit variance is symmetric about zero.



x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
0·00	0·5000	0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·97725
·01	·5040	·41	·6591	·81	·7910	·21	·8869	·61	·9463	·01	·97778
·02	·5080	·42	·6628	·82	·7939	·22	·8888	·62	·9474	·02	·97831
·03	·5120	·43	·6664	·83	·7967	·23	·8907	·63	·9484	·03	·97882
·04	·5160	·44	·6700	·84	·7995	·24	·8925	·64	·9495	·04	·97932
0·05	0·5199	0·45	0·6736	0·85	0·8023	1·25	0·8944	1·65	0·9505	2·05	0·97982
·06	·5239	·46	·6772	·86	·8051	·26	·8962	·66	·9515	·06	·98030
·07	·5279	·47	·6808	·87	·8078	·27	·8980	·67	·9525	·07	·98077
·08	·5319	·48	·6844	·88	·8106	·28	·8997	·68	·9535	·08	·98124
·09	·5359	·49	·6879	·89	·8133	·29	·9015	·69	·9545	·09	·98169
0·10	0·5398	0·50	0·6915	0·90	0·8159	1·30	0·9032	1·70	0·9554	2·10	0·98214
·11	·5438	·51	·6950	·91	·8186	·31	·9049	·71	·9564	·11	·98257
·12	·5478	·52	·6985	·92	·8212	·32	·9066	·72	·9573	·12	·98300
·13	·5517	·53	·7019	·93	·8238	·33	·9082	·73	·9582	·13	·98341
·14	·5557	·54	·7054	·94	·8264	·34	·9099	·74	·9591	·14	·98382
0·15	0·5596	0·55	0·7088	0·95	0·8289	1·35	0·9115	1·75	0·9599	2·15	0·98422
·16	·5636	·56	·7123	·96	·8315	·36	·9131	·76	·9608	·16	·98461
·17	·5675	·57	·7157	·97	·8340	·37	·9147	·77	·9616	·17	·98500
·18	·5714	·58	·7190	·98	·8365	·38	·9162	·78	·9625	·18	·98537
·19	·5753	·59	·7224	·99	·8389	·39	·9177	·79	·9633	·19	·98574
0·20	0·5793	0·60	0·7257	1·00	0·8413	1·40	0·9192	1·80	0·9641	2·20	0·98610
·21	·5832	·61	·7291	·01	·8438	·41	·9207	·81	·9649	·21	·98645
·22	·5871	·62	·7324	·02	·8461	·42	·9222	·82	·9656	·22	·98679
·23	·5910	·63	·7357	·03	·8485	·43	·9236	·83	·9664	·23	·98713
·24	·5948	·64	·7389	·04	·8508	·44	·9251	·84	·9671	·24	·98745
0·25	0·5987	0·65	0·7422	1·05	0·8531	1·45	0·9265	1·85	0·9678	2·25	0·98778
·26	·6026	·66	·7454	·06	·8554	·46	·9279	·86	·9686	·26	·98809
·27	·6064	·67	·7486	·07	·8577	·47	·9292	·87	·9693	·27	·98840
·28	·6103	·68	·7517	·08	·8599	·48	·9306	·88	·9699	·28	·98870
·29	·6141	·69	·7549	·09	·8621	·49	·9319	·89	·9706	·29	·98899
0·30	0·6179	0·70	0·7580	1·10	0·8643	1·50	0·9332	1·90	0·9713	2·30	0·98928
·31	·6217	·71	·7611	·11	·8665	·51	·9345	·91	·9719	·31	·98956
·32	·6255	·72	·7642	·12	·8686	·52	·9357	·92	·9726	·32	·98983
·33	·6293	·73	·7673	·13	·8708	·53	·9370	·93	·9732	·33	·99010
·34	·6331	·74	·7704	·14	·8729	·54	·9382	·94	·9738	·34	·99036
0·35	0·6368	0·75	0·7734	1·15	0·8749	1·55	0·9394	1·95	0·9744	2·35	0·99061
·36	·6406	·76	·7764	·16	·8770	·56	·9406	·96	·9750	·36	·99086
·37	·6443	·77	·7794	·17	·8790	·57	·9418	·97	·9756	·37	·99111
·38	·6480	·78	·7823	·18	·8810	·58	·9429	·98	·9761	·38	·99134
·39	·6517	·79	·7852	·19	·8830	·59	·9441	·99	·9767	·39	·99158
0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·9772	2·40	0·99180

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
2·40	0·99180	2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918
·41	·99202	·56	·99477	·71	·99664	·86	·99788	·01	·99869	·16	·99921
·42	·99224	·57	·99492	·72	·99674	·87	·99795	·02	·99874	·17	·99924
·43	·99245	·58	·99506	·73	·99683	·88	·99801	·03	·99878	·18	·99926
·44	·99266	·59	·99520	·74	·99693	·89	·99807	·04	·99882	·19	·99929
2·45	0·99286	2·60	0·99534	2·75	0·99702	2·90	0·99813	3·05	0·99886	3·20	0·99931
·46	·99305	·61	·99547	·76	·99711	·91	·99819	·06	·99889	·21	·99934
·47	·99324	·62	·99560	·77	·99720	·92	·99825	·07	·99893	·22	·99936
·48	·99343	·63	·99573	·78	·99728	·93	·99831	·08	·99896	·23	·99938
·49	·99361	·64	·99585	·79	·99736	·94	·99836	·09	·99900	·24	·99940
2·50	0·99379	2·65	0·99598	2·80	0·99744	2·95	0·99841	3·10	0·99903	3·25	0·99942
·51	·99396	·66	·99609	·81	·99752	·96	·99846	·11	·99906	·26	·99944
·52	·99413	·67	·99621	·82	·99760	·97	·99851	·12	·99910	·27	·99946
·53	·99430	·68	·99632	·83	·99767	·98	·99856	·13	·99913	·28	·99948
·54	·99446	·69	·99643	·84	·99774	·99	·99861	·14	·99916	·29	·99950
2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918	3·30	0·99952

The critical table below gives on the left the range of values of x for which $\Phi(x)$ takes the value on the right, correct to the last figure given; in critical cases, take the upper of the two values of $\Phi(x)$ indicated.

3·075	0·9990	3·263	0·9994	3·731	0·99990	3·916	0·99995
3·105	0·9991	3·320	0·9995	3·759	0·99991	3·976	0·99996
3·138	0·9992	3·389	0·9996	3·791	0·99992	4·055	0·99997
3·174	0·9993	3·480	0·9997	3·826	0·99993	4·173	0·99998
3·215	0·9994	3·615	0·9998	3·867	0·99994	4·417	0·99999
			0·9999		0·99995		1·00000

When $x > 3{\cdot}3$ the formula $1-\Phi(x) \doteqdot \frac{e^{-\frac{1}{2}x^2}}{x\sqrt{2\pi}}\left[1-\frac{1}{x^2}+\frac{3}{x^4}-\frac{15}{x^6}+\frac{105}{x^8}\right]$ is very accurate, with relative error less than $945/x^{10}$.

TABLE 5. PERCENTAGE POINTS OF THE NORMAL DISTRIBUTION

This table gives percentage points $x(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{2\pi}}\int_{x(P)}^{\infty} e^{-\frac{1}{2}t^2}\,dt.$$

If X is a variable, normally distributed with zero mean and unit variance, $P/100$ is the probability that $X \geqslant x(P)$. The lower P per cent points are given by symmetry as $-x(P)$, and the probability that $|X| \geqslant x(P)$ is $2P/100$.



P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$
50	0·0000	5·0	1·6449	3·0	1·8808	2·0	2·0537	1·0	2·3263	0·10	3·0902
45	0·1257	4·8	1·6646	2·9	1·8957	1·9	2·0749	0·9	2·3656	0·09	3·1214
40	0·2533	4·6	1·6849	2·8	1·9110	1·8	2·0969	0·8	2·4089	0·08	3·1559
35	0·3853	4·4	1·7060	2·7	1·9268	1·7	2·1201	0·7	2·4573	0·07	3·1947
30	0·5244	4·2	1·7279	2·6	1·9431	1·6	2·1444	0·6	2·5121	0·06	3·2389
25	0·6745	4·0	1·7507	2·5	1·9600	1·5	2·1701	0·5	2·5758	0·05	3·2905
20	0·8416	3·8	1·7744	2·4	1·9774	1·4	2·1973	0·4	2·6521	0·01	3·7190
15	1·0364	3·6	1·7991	2·3	1·9954	1·3	2·2262	0·3	2·7478	0·005	3·8906
10	1·2816	3·4	1·8250	2·2	2·0141	1·2	2·2571	0·2	2·8782	0·001	4·2649
5	1·6449	3·2	1·8522	2·1	2·0335	1·1	2·2904	0·1	3·0902	0·0005	4·4172

TABLE 10. PERCENTAGE POINTS OF THE t-DISTRIBUTION

This table gives percentage points $t_\nu(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{\nu\pi}} \frac{\Gamma(\frac{1}{2}\nu+\frac{1}{2})}{\Gamma(\frac{1}{2}\nu)} \int_{t_\nu(P)}^{\infty} \frac{dt}{(1+t^2/\nu)^{\frac{1}{2}(\nu+1)}}.$$

Let X_1 and X_2 be independent random variables having a normal distribution with zero mean and unit variance and a χ^2-distribution with ν degrees of freedom respectively; then $t = X_1/\sqrt{X_2/\nu}$ has Student's t-distribution with ν degrees of freedom, and the probability that $t \geqslant t_\nu(P)$ is $P/100$. The lower percentage points are given by symmetry as $-t_\nu(P)$, and the probability that $|t| \geqslant t_\nu(P)$ is $2P/100$.



The limiting distribution of t as ν tends to infinity is the normal distribution with zero mean and unit variance. When ν is large interpolation in ν should be harmonic.

P	40	30	25	20	15	10	5	2·5	1	0·5	0·1	0·05
$\nu = 1$	0·3249	0·7265	1·0000	1·3764	1·963	3·078	6·314	12·71	31·82	63·66	318·3	636·6
2	0·2887	0·6172	0·8165	1·0607	1·386	1·886	2·920	4·303	6·965	9·925	22·33	31·60
3	0·2767	0·5844	0·7649	0·9785	1·250	1·638	2·353	3·182	4·541	5·841	10·21	12·92
4	0·2707	0·5686	0·7407	0·9410	1·190	1·533	2·132	2·776	3·747	4·604	7·173	8·610
5	0·2672	0·5594	0·7267	0·9195	1·156	1·476	2·015	2·571	3·365	4·032	5·893	6·869
6	0·2648	0·5534	0·7176	0·9057	1·134	1·440	1·943	2·447	3·143	3·707	5·208	5·959
7	0·2632	0·5491	0·7111	0·8960	1·119	1·415	1·895	2·365	2·998	3·499	4·785	5·408
8	0·2619	0·5459	0·7064	0·8889	1·108	1·397	1·860	2·306	2·896	3·355	4·501	5·041
9	0·2610	0·5435	0·7027	0·8834	1·100	1·383	1·833	2·262	2·821	3·250	4·297	4·781
10	0·2602	0·5415	0·6998	0·8791	1·093	1·372	1·812	2·228	2·764	3·169	4·144	4·587
11	0·2596	0·5399	0·6974	0·8755	1·088	1·363	1·796	2·201	2·718	3·106	4·025	4·437
12	0·2590	0·5386	0·6955	0·8726	1·083	1·356	1·782	2·179	2·681	3·055	3·930	4·318
13	0·2586	0·5375	0·6938	0·8702	1·079	1·350	1·771	2·160	2·650	3·012	3·852	4·221
14	0·2582	0·5366	0·6924	0·8681	1·076	1·345	1·761	2·145	2·624	2·977	3·787	4·140
15	0·2579	0·5357	0·6912	0·8662	1·074	1·341	1·753	2·131	2·602	2·947	3·733	4·073
16	0·2576	0·5350	0·6901	0·8647	1·071	1·337	1·746	2·120	2·583	2·921	3·686	4·015
17	0·2573	0·5344	0·6892	0·8633	1·069	1·333	1·740	2·110	2·567	2·898	3·646	3·965
18	0·2571	0·5338	0·6884	0·8620	1·067	1·330	1·734	2·101	2·552	2·878	3·610	3·922
19	0·2569	0·5333	0·6876	0·8610	1·066	1·328	1·729	2·093	2·539	2·861	3·579	3·883
20	0·2567	0·5329	0·6870	0·8600	1·064	1·325	1·725	2·086	2·528	2·845	3·552	3·850
21	0·2566	0·5325	0·6864	0·8591	1·063	1·323	1·721	2·080	2·518	2·831	3·527	3·819
22	0·2564	0·5321	0·6858	0·8583	1·061	1·321	1·717	2·074	2·508	2·819	3·505	3·792
23	0·2563	0·5317	0·6853	0·8575	1·060	1·319	1·714	2·069	2·500	2·807	3·485	3·768
24	0·2562	0·5314	0·6848	0·8569	1·059	1·318	1·711	2·064	2·492	2·797	3·467	3·745
25	0·2561	0·5312	0·6844	0·8562	1·058	1·316	1·708	2·060	2·485	2·787	3·450	3·725
26	0·2560	0·5309	0·6840	0·8557	1·058	1·315	1·706	2·056	2·479	2·779	3·435	3·707
27	0·2559	0·5306	0·6837	0·8551	1·057	1·314	1·703	2·052	2·473	2·771	3·421	3·690
28	0·2558	0·5304	0·6834	0·8546	1·056	1·313	1·701	2·048	2·467	2·763	3·408	3·674
29	0·2557	0·5302	0·6830	0·8542	1·055	1·311	1·699	2·045	2·462	2·756	3·396	3·659
30	0·2556	0·5300	0·6828	0·8538	1·055	1·310	1·697	2·042	2·457	2·750	3·385	3·646
32	0·2555	0·5297	0·6822	0·8530	1·054	1·309	1·694	2·037	2·449	2·738	3·365	3·622
34	0·2553	0·5294	0·6818	0·8523	1·052	1·307	1·691	2·032	2·441	2·728	3·348	3·601
36	0·2552	0·5291	0·6814	0·8517	1·052	1·306	1·688	2·028	2·434	2·719	3·333	3·582
38	0·2551	0·5288	0·6810	0·8512	1·051	1·304	1·686	2·024	2·429	2·712	3·319	3·566
40	0·2550	0·5286	0·6807	0·8507	1·050	1·303	1·684	2·021	2·423	2·704	3·307	3·551
50	0·2547	0·5278	0·6794	0·8489	1·047	1·299	1·676	2·009	2·403	2·678	3·261	3·496
60	0·2545	0·5272	0·6786	0·8477	1·045	1·296	1·671	2·000	2·390	2·660	3·232	3·460
120	0·2539	0·5258	0·6765	0·8446	1·041	1·289	1·658	1·980	2·358	2·617	3·160	3·373
∞	0·2533	0·5244	0·6745	0·8416	1·036	1·282	1·645	1·960	2·326	2·576	3·090	3·291

TABLE 12(*a*). 10 PER CENT POINTS OF THE *F*-DISTRIBUTION

The function tabulated is $F(P) = F(P|\nu_1, \nu_2)$ defined by the equation

$$\frac{P}{100} = \frac{\Gamma(\frac{1}{2}\nu_1 + \frac{1}{2}\nu_2)}{\Gamma(\frac{1}{2}\nu_1)\,\Gamma(\frac{1}{2}\nu_2)}\,\nu_1^{\frac{1}{2}\nu_1}\,\nu_2^{\frac{1}{2}\nu_2}\int_{F(P)}^{\infty}\frac{F^{\frac{1}{2}\nu_1-1}}{(\nu_2+\nu_1 F)^{\frac{1}{2}(\nu_1+\nu_2)}}\,dF,$$

for P = 10, 5, 2·5, 1, 0·5 and 0·1. The lower percentage points, that is the values $F'(P) = F'(P|\nu_1, \nu_2)$ such that the probability that $F \leqslant F'(P)$ is equal to $P/100$, may be found by the formula

$$F'(P|\nu_1, \nu_2) = 1/F(P|\nu_2, \nu_1).$$



(This shape applies only when $\nu_1 \geqslant 3$. When $\nu_1 < 3$ the mode is at the origin.)

ν_1 =	1	2	3	4	5	6	7	8	10	12	24	∞
ν_2 = 1	39·86	49·50	53·59	55·83	57·24	58·20	58·91	59·44	60·19	60·71	62·00	63·33
2	8·526	9·000	9·162	9·243	9·293	9·326	9·349	9·367	9·392	9·408	9·450	9·491
3	5·538	5·462	5·391	5·343	5·309	5·285	5·266	5·252	5·230	5·216	5·176	5·134
4	4·545	4·325	4·191	4·107	4·051	4·010	3·979	3·955	3·920	3·896	3·831	3·761
5	4·060	3·780	3·619	3·520	3·453	3·405	3·368	3·339	3·297	3·268	3·191	3·105
6	3·776	3·463	3·289	3·181	3·108	3·055	3·014	2·983	2·937	2·905	2·818	2·722
7	3·589	3·257	3·074	2·961	2·883	2·827	2·785	2·752	2·703	2·668	2·575	2·471
8	3·458	3·113	2·924	2·806	2·726	2·668	2·624	2·589	2·538	2·502	2·404	2·293
9	3·360	3·006	2·813	2·693	2·611	2·551	2·505	2·469	2·416	2·379	2·277	2·159
10	3·285	2·924	2·728	2·605	2·522	2·461	2·414	2·377	2·323	2·284	2·178	2·055
11	3·225	2·860	2·660	2·536	2·451	2·389	2·342	2·304	2·248	2·209	2·100	1·972
12	3·177	2·807	2·606	2·480	2·394	2·331	2·283	2·245	2·188	2·147	2·036	1·904
13	3·136	2·763	2·560	2·434	2·347	2·283	2·234	2·195	2·138	2·097	1·983	1·846
14	3·102	2·726	2·522	2·395	2·307	2·243	2·193	2·154	2·095	2·054	1·938	1·797
15	3·073	2·695	2·490	2·361	2·273	2·208	2·158	2·119	2·059	2·017	1·899	1·755
16	3·048	2·668	2·462	2·333	2·244	2·178	2·128	2·088	2·028	1·985	1·866	1·718
17	3·026	2·645	2·437	2·308	2·218	2·152	2·102	2·061	2·001	1·958	1·836	1·686
18	3·007	2·624	2·416	2·286	2·196	2·130	2·079	2·038	1·977	1·933	1·810	1·657
19	2·990	2·606	2·397	2·266	2·176	2·109	2·058	2·017	1·956	1·912	1·787	1·631
20	2·975	2·589	2·380	2·249	2·158	2·091	2·040	1·999	1·937	1·892	1·767	1·607
21	2·961	2·575	2·365	2·233	2·142	2·075	2·023	1·982	1·920	1·875	1·748	1·586
22	2·949	2·561	2·351	2·219	2·128	2·060	2·008	1·967	1·904	1·859	1·731	1·567
23	2·937	2·549	2·339	2·207	2·115	2·047	1·995	1·953	1·890	1·845	1·716	1·549
24	2·927	2·538	2·327	2·195	2·103	2·035	1·983	1·941	1·877	1·832	1·702	1·533
25	2·918	2·528	2·317	2·184	2·092	2·024	1·971	1·929	1·866	1·820	1·689	1·518
26	2·909	2·519	2·307	2·174	2·082	2·014	1·961	1·919	1·855	1·809	1·677	1·504
27	2·901	2·511	2·299	2·165	2·073	2·005	1·952	1·909	1·845	1·799	1·666	1·491
28	2·894	2·503	2·291	2·157	2·064	1·996	1·943	1·900	1·836	1·790	1·656	1·478
29	2·887	2·495	2·283	2·149	2·057	1·988	1·935	1·892	1·827	1·781	1·647	1·467
30	2·881	2·489	2·276	2·142	2·049	1·980	1·927	1·884	1·819	1·773	1·638	1·456
32	2·869	2·477	2·263	2·129	2·036	1·967	1·913	1·870	1·805	1·758	1·622	1·437
34	2·859	2·466	2·252	2·118	2·024	1·955	1·901	1·858	1·793	1·745	1·608	1·419
36	2·850	2·456	2·243	2·108	2·014	1·945	1·891	1·847	1·781	1·734	1·595	1·404
38	2·842	2·448	2·234	2·099	2·005	1·935	1·881	1·838	1·772	1·724	1·584	1·390
40	2·835	2·440	2·226	2·091	1·997	1·927	1·873	1·829	1·763	1·715	1·574	1·377
60	2·791	2·393	2·177	2·041	1·946	1·875	1·819	1·775	1·707	1·657	1·511	1·291
120	2·748	2·347	2·130	1·992	1·896	1·824	1·767	1·722	1·652	1·601	1·447	1·193
∞	2·706	2·303	2·084	1·945	1·847	1·774	1·717	1·670	1·599	1·546	1·383	1·000

TABLE 12(*b*). 5 PER CENT POINTS OF THE *F*-DISTRIBUTION

If $F = \frac{X_1}{\nu_1} \Big/ \frac{X_2}{\nu_2}$, where X_1 and X_2 are independent random variables distributed as χ^2 with ν_1 and ν_2 degrees of freedom respectively, then the probabilities that $F \geqslant F(P)$ and that $F \leqslant F'(P)$ are both equal to $P/100$. Linear interpolation in ν_1 and ν_2 will generally be sufficiently accurate except when either $\nu_1 > 12$ or $\nu_2 > 40$, when harmonic interpolation should be used.



(This shape applies only when $\nu_1 \geqslant 3$. When $\nu_1 < 3$ the mode is at the origin.)

$\nu_1 =$	1	2	3	4	5	6	7	8	10	12	24	∞
$\nu_2 = 1$	161·4	199·5	215·7	224·6	230·2	234·0	236·8	238·9	241·9	243·9	249·1	254·3
2	18·51	19·00	19·16	19·25	19·30	19·33	19·35	19·37	19·40	19·41	19·45	19·50
3	10·13	9·552	9·277	9·117	9·013	8·941	8·887	8·845	8·786	8·745	8·639	8·526
4	7·709	6·944	6·591	6·388	6·256	6·163	6·094	6·041	5·964	5·912	5·774	5·628
5	6·608	5·786	5·409	5·192	5·050	4·950	4·876	4·818	4·735	4·678	4·527	4·365
6	5·987	5·143	4·757	4·534	4·387	4·284	4·207	4·147	4·060	4·000	3·841	3·669
7	5·591	4·737	4·347	4·120	3·972	3·866	3·787	3·726	3·637	3·575	3·410	3·230
8	5·318	4·459	4·066	3·838	3·687	3·581	3·500	3·438	3·347	3·284	3·115	2·928
9	5·117	4·256	3·863	3·633	3·482	3·374	3·293	3·230	3·137	3·073	2·900	2·707
10	4·965	4·103	3·708	3·478	3·326	3·217	3·135	3·072	2·978	2·913	2·737	2·538
11	4·844	3·982	3·587	3·357	3·204	3·095	3·012	2·948	2·854	2·788	2·609	2·404
12	4·747	3·885	3·490	3·259	3·106	2·996	2·913	2·849	2·753	2·687	2·505	2·296
13	4·667	3·806	3·411	3·179	3·025	2·915	2·832	2·767	2·671	2·604	2·420	2·206
14	4·600	3·739	3·344	3·112	2·958	2·848	2·764	2·699	2·602	2·534	2·349	2·131
15	4·543	3·682	3·287	3·056	2·901	2·790	2·707	2·641	2·544	2·475	2·288	2·066
16	4·494	3·634	3·239	3·007	2·852	2·741	2·657	2·591	2·494	2·425	2·235	2·010
17	4·451	3·592	3·197	2·965	2·810	2·699	2·614	2·548	2·450	2·381	2·190	1·960
18	4·414	3·555	3·160	2·928	2·773	2·661	2·577	2·510	2·412	2·342	2·150	1·917
19	4·381	3·522	3·127	2·895	2·740	2·628	2·544	2·477	2·378	2·308	2·114	1·878
20	4·351	3·493	3·098	2·866	2·711	2·599	2·514	2·447	2·348	2·278	2·082	1·843
21	4·325	3·467	3·072	2·840	2·685	2·573	2·488	2·420	2·321	2·250	2·054	1·812
22	4·301	3·443	3·049	2·817	2·661	2·549	2·464	2·397	2·297	2·226	2·028	1·783
23	4·279	3·422	3·028	2·796	2·640	2·528	2·442	2·375	2·275	2·204	2·005	1·757
24	4·260	3·403	3·009	2·776	2·621	2·508	2·423	2·355	2·255	2·183	1·984	1·733
25	4·242	3·385	2·991	2·759	2·603	2·490	2·405	2·337	2·236	2·165	1·964	1·711
26	4·225	3·369	2·975	2·743	2·587	2·474	2·388	2·321	2·220	2·148	1·946	1·691
27	4·210	3·354	2·960	2·728	2·572	2·459	2·373	2·305	2·204	2·132	1·930	1·672
28	4·196	3·340	2·947	2·714	2·558	2·445	2·359	2·291	2·190	2·118	1·915	1·654
29	4·183	3·328	2·934	2·701	2·545	2·432	2·346	2·278	2·177	2·104	1·901	1·638
30	4·171	3·316	2·922	2·690	2·534	2·421	2·334	2·266	2·165	2·092	1·887	1·622
32	4·149	3·295	2·901	2·668	2·512	2·399	2·313	2·244	2·142	2·070	1·864	1·594
34	4·130	3·276	2·883	2·650	2·494	2·380	2·294	2·225	2·123	2·050	1·843	1·569
36	4·113	3·259	2·866	2·634	2·477	2·364	2·277	2·209	2·106	2·033	1·824	1·547
38	4·098	3·245	2·852	2·619	2·463	2·349	2·262	2·194	2·091	2·017	1·808	1·527
40	4·085	3·232	2·839	2·606	2·449	2·336	2·249	2·180	2·077	2·003	1·793	1·509
60	4·001	3·150	2·758	2·525	2·368	2·254	2·167	2·097	1·993	1·917	1·700	1·389
120	3·920	3·072	2·680	2·447	2·290	2·175	2·087	2·016	1·910	1·834	1·608	1·254
∞	3·841	2·996	2·605	2·372	2·214	2·099	2·010	1·938	1·831	1·752	1·517	1·000

Tables of the Studentized Range, alpha= 0.10

Denominator	Numerator											
df	2	3	4	5	6	7	8	9	10	11	12	13
1	8.93	13.44	16.36	18.49	20.15	21.50	22.64	23.62	24.48	25.24	25.92	26.54
2	4.13	5.73	6.77	7.54	8.14	8.63	9.05	9.41	9.73	10.01	10.26	10.49
3	3.33	4.47	5.20	5.74	6.16	6.51	6.81	7.06	7.29	7.49	7.67	7.83
4	3.02	3.98	4.59	5.04	5.39	5.68	5.93	6.14	6.33	6.49	6.65	6.78
5	2.85	3.72	4.26	4.66	4.98	5.24	5.46	5.65	5.82	5.97	6.10	6.22
6	2.75	3.56	4.07	4.44	4.73	4.97	5.17	5.34	5.50	5.64	5.76	5.88
7	2.68	3.45	3.93	4.28	4.56	4.78	4.97	5.14	5.28	5.41	5.53	5.64
8	2.63	3.37	3.83	4.17	4.43	4.65	4.83	4.99	5.13	5.25	5.36	5.46
9	2.59	3.32	3.76	4.08	4.34	4.55	4.72	4.87	5.01	5.13	5.23	5.33
10	2.56	3.27	3.70	4.02	4.26	4.47	4.64	4.78	4.91	5.03	5.13	5.23
11	2.54	3.23	3.66	3.97	4.21	4.40	4.57	4.71	4.84	4.95	5.05	5.15
12	2.52	3.20	3.62	3.92	4.16	4.35	4.51	4.65	4.78	4.89	4.99	5.08
13	2.50	3.18	3.59	3.89	4.12	4.30	4.46	4.60	4.72	4.83	4.93	5.02
14	2.49	3.16	3.56	3.85	4.08	4.27	4.42	4.56	4.68	4.79	4.88	4.97
15	2.48	3.14	3.54	3.83	4.05	4.24	4.39	4.52	4.64	4.75	4.84	4.93
16	2.47	3.12	3.52	3.80	4.03	4.21	4.36	4.49	4.61	4.71	4.81	4.89
17	2.46	3.11	3.50	3.78	4.00	4.18	4.33	4.46	4.58	4.68	4.77	4.86
18	2.45	3.10	3.49	3.77	3.98	4.16	4.31	4.44	4.55	4.65	4.75	4.83
19	2.45	3.09	3.47	3.75	3.97	4.14	4.29	4.42	4.53	4.63	4.72	4.80
20	2.44	3.08	3.46	3.74	3.95	4.12	4.27	4.40	4.51	4.61	4.70	4.78
21	2.43	3.07	3.45	3.72	3	4.11	4.26	4.38	4.49	4.59	4.68	4.76
22	2.43	3.06	3.44	3.71	3.92	4.10	4.24	4.36	4.47	4.57	4.66	4.74
23	2.42	3.05	3.43	3.70	3.91	4.08	4.23	4.35	4.46	4.56	4.64	4.72
24	2.42	3.05	3.42	3.69	3.90	4.07	4.21	4.34	4.45	4.54	4.63	4.71
25	2.42	3.04	3.42	3.68	3.89	4.06	4.20	4.32	4.43	4.53	4.61	4.69
26	2.41	3.04	3.41	3.68	3.88	4.05	4.19	4.31	4.42	4.52	4.60	4.68
27	2.41	3.03	3.40	3.67	3.87	4.04	4.18	4.30	4.41	4.50	4.59	4.67
28	2.41	3.03	3.40	3.66	3.87	4.03	4.17	4.29	4.40	4.49	4.58	4.66
29	2.40	3.02	3.39	3.65	3.86	4.02	4.16	4.28	4.39	4.48	4.57	4.65
30	2.40	3.02	3.39	3.65	3.85	4.02	4.16	4.28	4.38	4.47	4.56	4.64
40	2.38	2.99	3.35	3.61	3.80	3.96	4.10	4.22	4.32	4.41	4.49	4.56
60	2.36	2.96	3.31	3.56	3.76	3.91	4.04	4.16	4.25	4.34	4.42	4.49
80	2.35	2.95	3.29	3.54	3.73	3.89	4.01	4.13	4.22	4.31	4.39	4.46
120	2.34	2.93	3.28	3.52	3.71	3.86	3.99	4.10	4.19	4.28	4.35	4.42
240	2.34	2.92	3.26	3.50	3.68	3.83	3.96	4.07	4.16	4.24	4.32	4.39
∞	2.33	2.90	3.24	3.48	3.66	3.81	3.93	4.04	4.13	4.21	4.29	4.35